

JOURNAL OF MATHEMATICAL ANALYSIS AND APPLICATIONS 127, 211–225 (1987)

## On Regular Extensions of Contents and Measures

WOLFGANG ADAMSKI

*Mathematisches Institut, Universität München,  
D-8000 Munich 2, West Germany**Submitted by Ky Fan*

Received December 30, 1985

### INTRODUCTION

Let  $\mathcal{A}$  be an algebra and  $\mathcal{K} \subset \mathcal{L}$  lattices of subsets of a set  $X$ . Furthermore, let  $\mu$  be a semifinite content on  $\mathcal{A}$  such that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ . Then our main result (3.4 and 3.6) states that  $\mu$  can be extended to a semifinite,  $\mathcal{L}$ -regular content defined on an algebra containing both  $\mathcal{A}$  and  $\mathcal{L}$ . If, in addition,  $\mu$  is a measure and  $\mathcal{L}$  is sequentially dominated by  $\mathcal{A}$ , then  $\mu$  can be extended to a semifinite,  $\mathcal{L}_\delta$ -regular measure defined on a  $\sigma$ -algebra containing both  $\mathcal{A}$  and  $\mathcal{L}$ . Since, in the special case  $\mathcal{K} \subset \mathcal{A}$ ,  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  iff  $\mu$  is  $\mathcal{K}$ -regular, our result is a common generalization of measure extension theorems due to Bachman and Sultan [5, 2.1], Lembcke [11, 3.1 and 4.5], Szeto [18, 2.2] and the author [3, 3.2 and 3.3(a)].

Our proceeding is based on the concept of a tight set function [2 and 3]. Some fundamental properties of tight set functions are compiled in Section 1. An essential tool for proving our main theorem is the fact that every supermodular set function defined on a lattice of sets admits a tight majorant. This result which is also important in cooperative game theory (see [6, 9, 16]) is proved in Section 2. A similar result concerning the existence of tight minorants for certain submodular set functions which has been proved in [1] is the basic tool for the derivation of another general measure extension theorem (3.12). As a consequence of this theorem, it is shown, among others, that every tight Baire measure on a [completely] Hausdorff space can be [uniquely] extended to a Radon measure (3.16).

## 1. DEFINITIONS AND AUXILIARY RESULTS

Let  $X$  be an arbitrary set. If  $A$  is a subset of  $X$  and  $f$  is a function defined on  $X$ , then  $f|A$  denotes the restriction of  $f$  on  $A$ . A *paving* (in  $X$ ) is a subset of  $\mathcal{P}(X)$ , the power set of  $X$ , which contains the empty set. A paving that is closed under finite [countable] intersections and finite unions is called a *lattice* [ $\delta$ -*lattice*] of sets.

Let  $\mathcal{C}$  be a paving in  $X$ . Then  $\alpha(\mathcal{C})$  [ $\sigma(\mathcal{C})$ ] denotes the  $[\sigma]$ -algebra generated by  $\mathcal{C}$ . Furthermore,  $\mathcal{C}_\delta$  denotes the family of all countable intersections of sets from  $\mathcal{C}$ .  $\mathcal{C}$  is said to be *semicompact* if every countable subfamily of  $\mathcal{C}$  having the finite intersection property has nonvoid intersection. In the following let  $\mathcal{K}$ ,  $\mathcal{L}$  be lattices of subsets of  $X$ .  $\mathcal{K}$  is said to be *sequentially dominated* by  $\mathcal{L}$  if whenever  $(K_n)$  is a sequence in  $\mathcal{K}$  with  $K_n \downarrow \emptyset$ , there exists a sequence  $(L_n)$  in  $\mathcal{L}$  such that  $L_n \downarrow \emptyset$  and  $K_n \subset L_n$  for all  $n \in \mathbb{N}$ . We say  $\mathcal{L}$  *separates*  $\mathcal{K}$  if whenever  $K_1, K_2 \in \mathcal{K}$  with  $K_1 \cap K_2 = \emptyset$ , there are disjoint  $\mathcal{L}$ -sets  $L_1, L_2$  such that  $K_i \subset L_i$  for  $i = 1, 2$ .

$\mathcal{F}(\mathcal{K}) := \{F \in \mathcal{P}(X) : F \cap K \in \mathcal{K} \text{ for all } K \in \mathcal{K}\}$  denotes the lattice of “local  $\mathcal{K}$ -sets.” Obviously  $X \in \mathcal{F}(\mathcal{K})$  and  $\mathcal{K} \subset \mathcal{F}(\mathcal{K}) \subset \mathcal{F}(\mathcal{K}_\delta)$ . Moreover, we have  $\mathcal{K} = \mathcal{F}(\mathcal{K})$  iff  $X \in \mathcal{K}$ .

Furthermore, we denote by  $N(\mathcal{K})$  the family of all  $[0, \infty]$ -valued set functions defined on  $\mathcal{K}$  and vanishing at  $\emptyset$ . If  $\mathcal{K}$  is an algebra, then a finitely [countably] additive element of  $N(\mathcal{K})$  is called a *content* [*measure*] on  $\mathcal{K}$ .

For a given  $\lambda \in N(\mathcal{K})$  we define set functions  $\lambda_*$ ,  $\lambda^*$  on  $\mathcal{P}(X)$  by  $\lambda_*(Q) := \sup\{\lambda(K) : K \in \mathcal{K}, K \subset Q\}$  and  $\lambda^*(Q) := \inf\{\lambda(K) : K \in \mathcal{K}, K \supset Q\}$  (with  $\inf \emptyset = \infty$ ). Moreover, we put  $\|\lambda\| := \lambda_*(X)$ .

Let  $\mathcal{C}$  be a subpaving of  $\mathcal{K}$ .  $\lambda \in N(\mathcal{K})$  is said to be

- (a)  *$\mathcal{C}$ -regular* if  $\lambda(K) = \sup\{\lambda(C) : C \in \mathcal{C}, C \subset K\}$  for  $K \in \mathcal{K}$ ;
- (b) *finite* if  $\lambda(K) < \infty$  for all  $K \in \mathcal{K}$ ;
- (c) *semifinite* if  $\lambda(K) = \infty$  implies  $\lambda(K) = \sup\{\lambda(K_0) : K_0 \in \mathcal{K}, K_0 \subset K, \lambda(K_0) < \infty\}$ ;
- (d) *supermodular* if  $\lambda(K_1) + \lambda(K_2) \leq \lambda(K_1 \cap K_2) + \lambda(K_1 \cup K_2)$  for  $K_1, K_2 \in \mathcal{K}$ ;
- (e) *modular* if  $\lambda(K_1) + \lambda(K_2) = \lambda(K_1 \cap K_2) + \lambda(K_1 \cup K_2)$  for  $K_1, K_2 \in \mathcal{K}$ ;
- (f) *tight* if  $\lambda(K_2) = \lambda(K_1) + \lambda_*(K_2 - K_1)$  for  $K_1, K_2 \in \mathcal{K}$  with  $K_1 \subset K_2$ ;
- (g)  *$\sigma$ -smooth at  $\emptyset$*  if  $\lim_n \lambda(K_n) = 0$  for every sequence  $(K_n)$  in  $\mathcal{K}$  satisfying  $K_n \downarrow \emptyset$  and  $\inf_n \lambda(K_n) < \infty$ .

For  $\rho \in N(\mathcal{P}(X))$  denote by  $\mathcal{M}(\rho) := \{A \in \mathcal{P}(X) : \rho(Q) = \rho(Q \cap A) + \rho(Q - A) \text{ for all } Q \subset X\}$  the family of so-called  $\rho$ -*measurable* subsets of  $X$ . It

is well known (see [10, Lemma 1]) that  $\mathcal{M}(\rho)$  is an algebra and  $\rho$  is a content on  $\mathcal{M}(\rho)$ . Furthermore, we need the following facts about tight set functions.

1.1. LEMMA [2, 2.1 and 2.2]. *Let  $\lambda \in N(\mathcal{X})$  be tight. Then*

- (a)  $\lambda$  is monotone and modular;
- (b)  $\mathcal{F}(\mathcal{X}) \subset \mathcal{M}(\lambda_*)$ ;
- (c)  $\lambda$  is  $\sigma$ -smooth at  $\emptyset$  if  $\mathcal{X}$  is semicompact.

1.2. LEMMA. *Let  $\lambda \in N(\mathcal{X})$  be tight with  $\|\lambda\| < \infty$ . Then we have  $\mathcal{M}(\lambda_*) = \{A \in \mathcal{P}(X) : \lambda_*(A) + \lambda_*(X-A) = \|\lambda\|\}$ .*

*Proof.* For  $A \in \mathcal{M}(\lambda_*)$ , the equality

$$\lambda_*(A) + \lambda_*(X-A) = \|\lambda\| \quad (1.1)$$

trivially holds. Now assume that  $A \in \mathcal{P}(X)$  satisfies (1.1). As  $\lambda_*$  is supermodular, we obtain, for any  $Q \subset X$ ,  $\lambda_*(Q) \geq \lambda_*(Q \cap A) + \lambda_*(Q - A) \geq \lambda_*(Q) + \lambda_*(A) - \lambda_*(Q \cup A) + \lambda_*(Q) + \lambda_*(X-A) - \lambda_*(Q \cup (X-A)) = \|\lambda\| + 2\lambda_*(Q) - \lambda_*(Q \cup A) - \lambda_*(Q \cup (X-A)) \geq \|\lambda\| + 2\lambda_*(Q) - \lambda_*(X) - \lambda_*(Q) = \lambda_*(Q)$  which implies  $A \in \mathcal{M}(\lambda_*)$ . ■

1.3. LEMMA. *Let  $\lambda \in N(\mathcal{X})$  be tight, semifinite, and  $\sigma$ -smooth at  $\emptyset$ . Define a set function  $\lambda^s \in N(\mathcal{P}(X))$  by*

$$\lambda^s(Q) := \sup\{\lambda^*(\bar{K}) : \bar{K} \in \mathcal{X}_\delta, \bar{K} \subset Q \text{ and } \lambda^*(\bar{K}) < \infty\} \text{ for } Q \in \mathcal{P}(X). \quad (1.2)$$

*Then we have*

- (a)  $\mathcal{M}(\lambda^s)$  is a  $\sigma$ -algebra containing both  $\mathcal{F}(\mathcal{X}_\delta)$  and  $\mathcal{M}(\lambda_*)$ .
- (b)  $\lambda^s \mid \mathcal{M}(\lambda^s)$  is a semifinite,  $\mathcal{X}_\delta$ -regular measure which extends  $\lambda$ .
- (c)  $\lambda_* \mid \mathcal{M}(\lambda_*) = \lambda^s \mid \mathcal{M}(\lambda_*)$ .
- (d) If  $\mathcal{X}$  is a  $\delta$ -lattice, then  $\lambda^s = \lambda_*$ .

*Proof.* See [2, 2.4–2.6], for a proof of (a), (b), and (d).

Ad (c): The inequality  $\lambda_* \leq \lambda^s$  is obvious. We first prove

$$\lambda_*(K \cap C) = \lambda^s(K \cap C) \quad \text{for } C \in \mathcal{M}(\lambda_*) \text{ and } K \in \mathcal{X} \text{ with } \lambda(K) < \infty. \quad (1.3)$$

Let  $C \in \mathcal{M}(\lambda_*)$  and  $K \in \mathcal{X}$  with  $\lambda(K) < \infty$  be given. By [2, 2.5], we then have  $\lambda^s(K \cap C) + \lambda^s(K - C) \leq \lambda^s(K) = \lambda(K) = \lambda_*(K \cap C) + \lambda_*(K - C) \leq \lambda^s(K \cap C) + \lambda^s(K - C)$ , hence  $\lambda^s(K \cap C) + \lambda^s(K - C) = \lambda_*(K \cap C) + \lambda_*(K - C) = \lambda(K) < \infty$  which, together with  $\lambda_* \leq \lambda^s$ , implies (1.3).

Now let  $C \in \mathcal{M}(\lambda_*)$  be given. Assume that  $\lambda_*(C) < \lambda^s(C)$ . As  $\lambda^s$  is  $\mathcal{K}_\delta$ -regular and semifinite, there is a  $\mathcal{K}_\delta$ -set  $\bar{K}$  with  $\bar{K} \subset C$  and  $\lambda_*(C) < \lambda^s(\bar{K}) = \lambda^*(\bar{K}) < \infty$ . Choose a decreasing sequence  $(K_n)$  in  $\mathcal{K}$  with  $\bar{K} = \bigcap_n K_n$  and  $\lambda(K_n) < \infty$  for all  $n$ . Then  $\bar{K} = \bigcap_n (K_n \cap C)$  and so, by (1.3),  $\lambda_*(C) < \lambda^s(\bar{K}) = \inf_n \lambda^s(K_n \cap C) = \inf_n \lambda_*(K_n \cap C) \leq \lambda_*(C)$ . This contradiction proves our claim. ■

Finally, we use the following topological notation. If  $X$  is a topological space, then any set of the form  $\{f=0\}$ , where  $f$  is a continuous real-valued function on  $X$  is called a *zero-set* in  $X$ . Complements of zero-sets are called *cozero-sets*. We denote by  $\mathcal{Z}(X)$ ,  $\mathcal{U}(X)$ ,  $\mathcal{F}(X)$ ,  $\mathcal{G}(X)$ , and  $\mathcal{K}(X)$  the family of all zero-, cozero-, closed, open, and compact sets in  $X$ , respectively.  $\mathcal{B}_0(X) := \sigma(\mathcal{Z}(X))$  [ $\mathcal{B}(X) := \sigma(\mathcal{F}(X))$ ] denotes the *Baire* [*Borel*]  $\sigma$ -algebra in  $X$ . Any measure defined on  $\mathcal{B}_0(X)$  [ $\mathcal{B}(X)$ ] is called a *Baire* [*Borel*] *measure* on  $X$ .

## 2. TIGHT MAJORANTS OF SUPERMODULAR SET FUNCTIONS

Throughout this section  $\mathcal{K}$  denotes a lattice of subsets of an arbitrary set  $X$ . We introduce a partial order in the set  $N(\mathcal{K})$  by defining  $\lambda_1 \leq \lambda_2$  iff  $\lambda_1(K) \leq \lambda_2(K)$  for all  $K \in \mathcal{K}$ .

**2.1. LEMMA.** *For  $a \in [0, \infty)$  let  $A^a := \{\lambda \in N(\mathcal{K}) : \lambda \text{ supermodular and } \|\lambda\| = a\}$ . Then*

(a) *every nonempty linearly ordered subset of  $A^a$  has an upper bound in  $A^a$  and*

(b) *every maximal element of  $A^a$  is tight.*

*Proof.* (a) Let  $A_0 := \{\lambda_i : i \in I\}$  be a nonvoid linearly ordered subset of  $A^a$ . Put  $\lambda(K) := \sup\{\lambda_i(K) : i \in I\}$  for  $K \in \mathcal{K}$ . It is clear that  $\lambda \in N(\mathcal{K})$  and  $\|\lambda\| = a$ . To prove the supermodularity of  $\lambda$ , let  $K_1, K_2 \in \mathcal{K}$  and  $\varepsilon > 0$  be given. Choose  $i_1, i_2$  in  $I$  such that  $\lambda(K_j) < \lambda_{i_j}(K_j) + \varepsilon/2$  for  $j = 1, 2$ . Suppose  $\lambda_{i_1} \leq \lambda_{i_2}$ . Then  $\lambda(K_1) + \lambda(K_2) < \lambda_{i_1}(K_1) + \lambda_{i_2}(K_2) + \varepsilon \leq \lambda_{i_2}(K_1) + \lambda_{i_2}(K_2) + \varepsilon \leq \lambda_{i_2}(K_1 \cap K_2) + \lambda_{i_2}(K_1 \cup K_2) + \varepsilon \leq \lambda(K_1 \cap K_2) + \lambda(K_1 \cup K_2) + \varepsilon$ . It follows that  $\lambda$  is supermodular. Hence  $\lambda \in A^a$  is an upper bound for  $A_0$ .

(b) Let  $\lambda$  be a maximal element of  $A^a$ . For any subset  $C$  of  $X$ , define a set function  $\lambda_C$  on  $\mathcal{K}$  by  $\lambda_C(K) := \lambda_*(K \cap C) + \lambda_*(K \cup C) - \lambda_*(C)$ ,  $K \in \mathcal{K}$ . We now prove that

$$\lambda_C \in A^a. \quad (2.1)$$

Trivially we have  $\lambda_C \in N(\mathcal{K})$ . In order to prove the supermodularity of  $\lambda_C$ , it suffices to show that, for any  $\mathcal{K}$ -sets  $K_1$  and  $K_2$ , we have  $\lambda_*(K_1 \cap C) +$

$\lambda_*(K_1 \cup C) + \lambda_*(K_2 \cap C) + \lambda_*(K_2 \cup C) \leq \lambda_*(K_1 \cap K_2 \cap C) + \lambda_*((K_1 \cap K_2) \cup C) + \lambda_*((K_1 \cup K_2) \cap C) + \lambda_*(K_1 \cup K_2 \cup C)$ . But this follows immediately from the two inequalities  $\lambda_*(K_1 \cap C) + \lambda_*(K_2 \cap C) \leq \lambda_*(K_1 \cap K_2 \cap C) + \lambda_*((K_1 \cup K_2) \cap C)$  and  $\lambda_*(K_1 \cup C) + \lambda_*(K_2 \cup C) \leq \lambda_*((K_1 \cap K_2) \cup C) + \lambda_*(K_1 \cup K_2 \cup C)$ . Furthermore, we have

$$\|\lambda_C\| = a. \quad (2.2)$$

It is clear that  $\lambda_C \geq \lambda \in A^a$  implies  $\|\lambda_C\| \geq a$ . Assuming  $\|\lambda_C\| > a$ , we have  $\lambda_*(K \cap C) + \lambda_*(K \cup C) - \lambda_*(C) > a$  for some  $\mathcal{K}$ -set  $K$  and hence  $\lambda_*(C) + a < \lambda_*(K \cap C) + \lambda_*(K \cup C) \leq \lambda_*(C) + a$ . This contradiction proves (2.2) and hence (2.1).

As we have  $\lambda_C \geq \lambda$ , the maximality of  $\lambda$  implies  $\lambda = \lambda_C$  for every subset  $C$  of  $X$ . In particular, for  $\mathcal{K}$ -sets  $K_1, K_2$  with  $K_1 \subset K_2$ , we obtain  $\lambda(K_1) = \lambda_{K_2 - K_1}(K_1) = \lambda(K_2) - \lambda_*(K_2 - K_1)$ . Hence  $\lambda$  is tight. ■

We can now prove the main result of this section. It is an essential tool for the derivation of a general measure extension theorem in the next section.

**2.2. THEOREM.** *For any supermodular  $\gamma \in N(\mathcal{K})$  there exists a tight  $\lambda \in N(\mathcal{K})$  such that  $\gamma \leq \lambda$  and  $\|\gamma\| = \|\lambda\|$ .*

*Proof.* If  $\|\gamma\| = \infty$ , then the tight set function  $\lambda$  defined by  $\lambda(K) = \infty$  for all nonvoid  $\mathcal{K}$ -sets  $K$  is appropriate. If  $a := \|\gamma\| < \infty$ , then, by 2.1(a) and Zorn's lemma, there is a maximal element  $\lambda \in A^a$  with  $\lambda \geq \gamma$ . Thus our claim follows from 2.1(b). ■

**2.3. Remarks.** (a) In the special case of a modular, monotone  $\gamma \in N(\mathcal{K})$  and  $X \in \mathcal{K}$ , 2.2 was proved independently by Pahl [14] and Szeto [18].

(b) If the supermodular set function  $\gamma$  is  $\{0, 1, \dots, n\}$ -valued (for some  $n \in N$ ), then a straightforward modification of the proof of 2.1 shows that the tight majorant  $\lambda$  in 2.2 can be chosen  $\{0, 1, \dots, n\}$ -valued, too.

(c) If  $\mathcal{K}$  is an algebra, then the tight set functions on  $\mathcal{K}$  are exactly the contents on  $\mathcal{K}$ . In this case, 2.2 is a well-known result ("core theorem") in cooperative game theory. The reader interested in the game theoretic aspects of nonnegative set functions is referred to [6, 9, 16].

(d) If, under the assumptions of 2.2,  $\gamma$  is finite but  $\|\gamma\| = \infty$ , then one cannot conclude that there is a tight majorant  $\lambda$  which is semifinite or even finite: Let  $\mathcal{K}$  be the family of all finite subsets of the real line and put  $\gamma(K) := (\text{card } K)^2$  for  $K \in \mathcal{K}$ . Then  $\gamma$  is supermodular and finite. An analysis of example (5.6) in [4] reveals that there is no modular  $\lambda \in N(\mathcal{K})$

satisfying  $\gamma \leq \lambda$  and  $\lambda(\{x\}) < \infty$  for all real numbers  $x$ . In particular, there does not exist any semifinite tight  $\lambda \in N(\mathcal{X})$  with  $\gamma \leq \lambda$ .

(e) If  $\gamma \in N(\mathcal{X})$  is supermodular and  $\sigma$ -smooth at  $\emptyset$ , then one cannot infer from 2.2 that there is a tight majorant  $\lambda$  which is also  $\sigma$ -smooth at  $\emptyset$ : Let  $X$  be any set with cardinality  $\aleph_1$ ,  $\mathcal{X}$  the power set of  $X$ , and  $\mathcal{D}$  the family of all subsets of  $X$  having a countable complement. For any  $K \in \mathcal{X}$ , put  $\gamma(K) = 1$  or  $0$  according as  $K \in \mathcal{D}$  or  $K \notin \mathcal{D}$ . Then  $\gamma \in N(\mathcal{X})$  is supermodular and  $\sigma$ -smooth at  $\emptyset$ . On the other hand, there is no probability measure  $\lambda$  on  $\mathcal{X}$  with  $\gamma \leq \lambda$ , since otherwise we would have  $\lambda(\{x\}) = 0$  for all  $x \in X$ , which is a contradiction to Ulam's theorem [13, Theorem 5.6].

### 3. REGULAR EXTENSIONS OF CONTENTS AND MEASURES

Throughout this section  $X$  denotes an arbitrary set, whereas  $\mathcal{X}$ ,  $\mathcal{L}$  are lattices of subsets of  $X$  and  $\mathcal{A}$  is an algebra in  $X$ . We begin with the following uniqueness statement.

3.1. PROPOSITION. Assume that  $\mathcal{A}$  separates  $\mathcal{X}$ .

(a) If  $\mathcal{A}'$  is an algebra containing  $\mathcal{A}$  and  $\mathcal{X}$ , then every finite content on  $\mathcal{A}$  has at most one extension to a  $\mathcal{X}$ -regular content on  $\mathcal{A}'$ .

(b) If  $\mathcal{A}'$  is a  $\sigma$ -algebra containing  $\mathcal{A}$  and  $\mathcal{X}$ , then every  $\sigma$ -finite measure on  $\mathcal{A}$  has at most one extension to a  $\mathcal{X}$ -regular measure on  $\mathcal{A}'$ .

*Proof.* (a) Let  $\mu$  be a finite content on  $\mathcal{A}$ . Suppose that  $\mu_1, \mu_2$  are  $\mathcal{X}$ -regular contents on  $\mathcal{A}'$  that extend  $\mu$ . In order to prove  $\mu_1 = \mu_2$  it suffices to show  $\mu_1(X - K) = \mu_2(X - K)$  for all  $K \in \mathcal{X}$ . Assume that  $\mu_1(X - K_0) \neq \mu_2(X - K_0)$ , say  $\mu_1(X - K_0) < \mu_2(X - K_0)$  for some  $\mathcal{X}$ -set  $K_0$ . Then there exists a  $\mathcal{X}$ -set  $K$  with  $K \subset X - K_0$  and  $\mu_1(X - K_0) < \mu_2(K)$ . Since  $\mathcal{A}$  separates  $\mathcal{X}$ , there are disjoint  $\mathcal{A}$ -sets  $A$  and  $A_0$  such that  $K \subset A$  and  $K_0 \subset A_0$ . This implies  $\mu(A) \leq \mu(X - A_0) \leq \mu_1(X - K_0) < \mu_2(K) \leq \mu(A)$  which is impossible.

(b) Let  $\mu$  be a  $\sigma$ -finite measure on  $\mathcal{A}$ . Then there is a disjoint sequence  $(X_n)$  in  $\mathcal{A}$  such that  $X = \bigcup_n X_n$  and  $\mu(X_n) < \infty$  for all  $n \in N$ . Let  $\nu$  and  $\tau$  be  $\mathcal{X}$ -regular measures on  $\mathcal{A}'$  which extend  $\mu$ . For  $n \in N$ ,  $A \in \mathcal{A}$ , and  $A' \in \mathcal{A}'$ , define  $\mu_n(A) := \mu(A \cap X_n)$ ,  $\nu_n(A') := \nu(A' \cap X_n)$ , and  $\tau_n(A') := \tau(A' \cap X_n)$ .  $\mu_n$  is a finite measure on  $\mathcal{A}$ , whereas  $\nu_n$  and  $\tau_n$  are finite,  $\mathcal{X}$ -regular measures on  $\mathcal{A}'$ . Since  $\nu_n$  and  $\tau_n$  extend  $\mu_n$ , we obtain  $\nu_n = \tau_n$ ,  $n \in N$ , by (a), and so  $\nu = \tau$ . ■

The example 3.5 in [3] shows that 3.1 does not remain valid for semifinite measures. The following extension result is an application of 2.2.

3.2. PROPOSITION. Let  $\gamma \in N(\mathcal{K})$  be supermodular with  $\|\gamma\| < \infty$ .

(a) If  $\gamma_* \mid \mathcal{A}$  is a content, then  $\gamma_* \mid \mathcal{A}$  can be extended to a  $\mathcal{K}$ -regular content on  $\alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{K}))$ .

(b) If  $\mathcal{K}$  is sequentially dominated by  $\mathcal{A}$  and  $\gamma_* \mid \mathcal{A}$  is a measure, then  $\gamma_* \mid \mathcal{A}$  can be extended to a  $\mathcal{K}_\delta$ -regular measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$ .

*Proof.* By 2.2, there is a tight  $\lambda \in N(\mathcal{K})$  such that  $\gamma \leq \lambda$  and  $\|\gamma\| = \|\lambda\|$ . Then, for any  $A \in \mathcal{A}$ , we have  $\gamma_*(A) \leq \lambda_*(A) \leq \|\lambda\| - \lambda_*(X - A) \leq \|\gamma\| - \gamma_*(X - A) = \gamma_*(A)$ , hence  $\gamma_*(A) = \lambda_*(A) = \|\lambda\| - \lambda_*(X - A)$  for all  $A \in \mathcal{A}$  which implies  $\mathcal{A} \subset \mathcal{M}(\lambda_*)$  by 1.2. On the other hand, we have  $\mathcal{F}(\mathcal{K}) \subset \mathcal{M}(\lambda_*)$  by 1.1(b). Thus  $\mathcal{A}' := \alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{K})) \subset \mathcal{M}(\lambda_*)$  which implies that  $\lambda_* \mid \mathcal{A}'$  is a  $\mathcal{K}$ -regular content that extends  $\gamma_* \mid \mathcal{A}$ . This proves (a).

Next we show that, under the additional assumptions of (b),  $\lambda$  is  $\sigma$ -smooth at  $\emptyset$ . For this purpose, let  $(K_n) \subset \mathcal{K}$  with  $K_n \downarrow \emptyset$ . Then there is a sequence  $(A_n)$  in  $\mathcal{A}$  such that  $A_n \downarrow \emptyset$  and  $K_n \subset A_n$  for all  $n$ . This implies  $\lambda(K_n) \leq \lambda_*(A_n) = \gamma_*(A_n) \rightarrow 0$  for  $n \rightarrow \infty$ . Hence  $\lambda$  is  $\sigma$ -smooth at  $\emptyset$ . Put  $\hat{\mathcal{A}} := \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$  and define  $\lambda^s$  by (1.2). Then we infer from 1.3 that  $\hat{\mathcal{A}} \subset \mathcal{M}(\lambda^s)$  and  $\lambda^s \mid \hat{\mathcal{A}}$  is a  $\mathcal{K}_\delta$ -regular measure which extends  $\gamma_* \mid \mathcal{A}$ . ■

Let  $\mu$  be a content defined on the algebra  $\mathcal{A}$ . Following the terminology of [15] we say that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  if  $\mu(A) = \sup\{\mu_*(K); K \in \mathcal{K}, K \subset A\}$  for all  $A \in \mathcal{A}$ .

3.3. Remarks. (a) If  $\mathcal{K} \subset \mathcal{A}$ , then  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  iff  $\mu$  is  $\mathcal{K}$ -regular.

(b) If  $\mu$  is finite, then  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  iff for every  $A \in \mathcal{A}$  and for every  $\varepsilon > 0$ , there exist sets  $A_0 \in \mathcal{A}$  and  $K \in \mathcal{K}$  such that  $A_0 \subset K \subset A$  and  $\mu(A - A_0) < \varepsilon$ . The latter condition goes back to Marczewski [12].

(c) If  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ , where  $\mathcal{K}$  is semicompact and  $\mu$  is semifinite, then  $\mu$  is  $\sigma$ -additive.

*Proof.* According to [2, 4.2(a)], it suffices to show that  $\mu$  is  $\sigma$ -smooth at  $\emptyset$ . But this can be done by standard arguments (see [12, (i), p. 118]). ■

We can now prove the following basic extension theorem.

3.4. THEOREM. Let  $\mu$  be a semifinite content on  $\mathcal{A}$  such that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ . Furthermore assume that  $\mathcal{K} \subset \mathcal{L}$ .

(a) Then  $\mu$  can be extended to a semifinite,  $\mathcal{L}$ -regular content on  $\alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{L}))$ . The extension is unique on every algebra  $\mathcal{A}'$  with  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{L}))$  if  $\mu$  is finite and  $\mathcal{A}$  separates  $\mathcal{L}$ .

(b) If, in addition,  $\mu$  is  $\sigma$ -additive and  $\mathcal{L}$  is sequentially dominated by  $\mathcal{A}$ , then  $\mu$  can be extended to a semifinite,  $\mathcal{L}_\delta$ -regular measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$ . The extension is unique on every  $\sigma$ -algebra  $\mathcal{A}'$  with  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$  if  $\mu$  is  $\sigma$ -finite and  $\mathcal{A}$  separates  $\mathcal{L}_\delta$ .

*Proof.* The uniqueness assertions follow directly from 3.1. Since  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ , so does  $\mathcal{L}$ . Thus in order to prove the existence statements it suffices to consider the special case  $\mathcal{K} = \mathcal{L}$ . We first assume  $\mu(X) < \infty$ . Putting  $\gamma(K) := \mu_*(K)$  for  $K \in \mathcal{K}$ , we have  $\mu = \gamma_* \upharpoonright \mathcal{A}$ . As  $\gamma \in N(\mathcal{K})$  is supermodular with  $\|\gamma\| = \mu(X) < \infty$ , our claim follows from 3.2. Now assume that  $\mu$  is semifinite. We only deal with the case where  $\mu$  is a semifinite measure on  $\mathcal{A}$  (and  $\mathcal{K}$  is sequentially dominated by  $\mathcal{A}$ ). The finitely additive case which is still simpler can be treated in an analogous way.

Let  $\mathcal{M}$  denote the family of all  $\mathcal{K}_\delta$ -regular measures  $\nu$  on  $\mathcal{A}' := \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$  satisfying  $\nu \upharpoonright \mathcal{A} \leq \mu$ . Obviously  $0 \in \mathcal{M}$ . We first prove that  $\mathcal{M}$  is inductively ordered. For this purpose let  $\mathcal{M}_0 \subset \mathcal{M}$  be linearly ordered and put  $\nu_0 := \sup \mathcal{M}_0$ , i.e.,  $\nu_0(A') = \sup \{\sum_n \nu_n(A_n) : (A_n) \subset \mathcal{A}' \text{ is a decomposition of } A' \text{ and } (\nu_n) \subset \mathcal{M}_0\}$ ,  $A' \in \mathcal{A}'$ . It is well known that  $\nu_0$  is a measure on  $\mathcal{A}'$  satisfying  $\nu_0 \geq \nu$  for all  $\nu \in \mathcal{M}_0$ . Furthermore,

$$\nu_0 \text{ is } \mathcal{K}_\delta\text{-regular.} \quad (3.1)$$

Let  $A' \in \mathcal{A}'$  and  $a < \nu_0(A')$ . Then there are measures  $\nu_1, \nu_2, \dots$  in  $\mathcal{M}_0$  and pairwise disjoint sets  $A_1, A_2, \dots$  in  $\mathcal{A}'$  such that  $A' = \bigcup_n A_n$  and  $a < \sum_n \nu_n(A_n)$ . Then  $a < \sum_{n=1}^k \nu_n(A_n)$  for some index  $k \in N$ . Choose a number  $k_0 \in \{1, \dots, k\}$  such that  $\nu_{k_0} \geq \nu_n$  for  $n = 1, \dots, k$ . Then  $a < \sum_{n=1}^k \nu_n(A_n) \leq \sum_{n=1}^k \nu_{k_0}(A_n) = \nu_{k_0}(\bigcup_{n=1}^k A_n) \leq \nu_{k_0}(A')$ . As  $\nu_{k_0}$  is  $\mathcal{K}_\delta$ -regular, there is a  $\mathcal{K}_\delta$ -set  $\bar{K} \subset A'$  such that  $a < \nu_{k_0}(\bar{K})$  and hence  $a < \nu_0(\bar{K})$ . This proves (3.1).

Now let  $A \in \mathcal{A}$  be given. Furthermore, let  $\nu_1, \nu_2, \dots \in \mathcal{M}_0$  and  $A_1, A_2, \dots$  pairwise disjoint  $\mathcal{A}'$ -sets with union  $A$ . For any  $k \in N$ , one can find an index  $k_0 \in \{1, \dots, k\}$  such that  $\nu_{k_0} \geq \nu_n$  for  $n = 1, \dots, k$ . This implies  $\sum_{n=1}^k \nu_n(A_n) \leq \sum_{n=1}^k \nu_{k_0}(A_n) = \nu_{k_0}(\bigcup_{n=1}^k A_n) \leq \nu_{k_0}(A) \leq \mu(A)$ . As  $k \in N$  was arbitrary, we obtain  $\sum_n \nu_n(A_n) \leq \mu(A)$  and hence  $\nu_0(A) \leq \mu(A)$ . Thus we have  $\nu_0 \upharpoonright \mathcal{A} \leq \mu$  which together with (3.1) implies  $\nu_0 \in \mathcal{M}$ . Hence  $\mathcal{M}$  is inductively ordered. By Zorn's lemma, there is a maximal element  $\tilde{\mu}$  in  $\mathcal{M}$ .

Now let  $A_0 \in \mathcal{A}$  with  $\mu(A_0) < \infty$  be given. Put  $\nu(A) := \mu(A \cap A_0) - \tilde{\mu}(A \cap A_0)$  for  $A \in \mathcal{A}$ . Then  $\nu$  is a finite measure on  $\mathcal{A}$ . Furthermore,

$$\mathcal{K} \text{ } \nu\text{-approximates } \mathcal{A}. \quad (3.2)$$

In order to prove (3.2), let  $A \in \mathcal{A}$  and  $\varepsilon > 0$  be given. There exist sets  $K \in \mathcal{K}$  and  $\tilde{A} \in \mathcal{A}$  such that  $\tilde{A} \subset K \subset A \cap A_0$  and  $\mu(A \cap A_0) < \mu(\tilde{A}) + \varepsilon$ . Then  $\nu(A) - \nu(\tilde{A}) = \mu(A \cap A_0) - \mu(\tilde{A}) - (\tilde{\mu}(A \cap A_0) - \tilde{\mu}(\tilde{A})) \leq \mu(A \cap A_0) - \mu(\tilde{A}) < \varepsilon$ . Thus (3.2) holds.



By the "finite case,"  $\nu$  can be extended to a  $\mathcal{K}_\delta$ -regular measure  $\tilde{\nu}$  on  $\mathcal{A}'$ . Then  $\tau := \tilde{\mu} + \tilde{\nu}$  is a  $\mathcal{K}_\delta$ -regular measure on  $\mathcal{A}'$ , and for any  $\mathcal{A}$ -set  $A$  we have  $\tau(A) = \tau(A \cap A_0) + \tau(A - A_0) = \tilde{\mu}(A \cap A_0) + \tilde{\nu}(A \cap A_0) + \tilde{\mu}(A - A_0) + \tilde{\nu}(A - A_0) = \tilde{\mu}(A \cap A_0) + \nu(A \cap A_0) + \tilde{\mu}(A - A_0) + \nu(A - A_0) = \mu(A \cap A_0) + \tilde{\mu}(A - A_0) \leq \mu(A \cap A_0) + \mu(A - A_0) = \mu(A)$ . Thus we have  $\tau \in \mathcal{M}$ . Since  $\tilde{\mu} \leq \tau$ , the maximality of  $\tilde{\mu}$  implies  $\tilde{\mu} = \tau$ . In particular, we have  $\infty > \mu(A_0) \geq \tilde{\mu}(A_0) = \tau(A_0) = \tilde{\mu}(A_0) + \tilde{\nu}(A_0)$ , hence  $\tilde{\nu}(A_0) = 0$  and so  $\mu(A_0) = \tilde{\mu}(A_0)$ . As the set  $A_0$  was arbitrary, we thus have proved

$$\mu(A) = \tilde{\mu}(A) \quad \text{for all } A \in \mathcal{A} \text{ with } \mu(A) < \infty. \quad (3.3)$$

For any  $A \in \mathcal{A}'$ , put  $\tilde{\mu}_1(A) := \sup\{\tilde{\mu}(B) : B \in \mathcal{A}', B \subset A \text{ and } \tilde{\mu}(B) < \infty\}$ . Then  $\tilde{\mu}_1$  is a semifinite measure on  $\mathcal{A}'$ . In addition,

$$\tilde{\mu}_1 \text{ is } \mathcal{K}_\delta\text{-regular.} \quad (3.4)$$

Let  $A \in \mathcal{A}'$  and  $a < \tilde{\mu}_1(A)$ . There exists a set  $B \in \mathcal{A}'$  with  $B \subset A$  and  $a < \tilde{\mu}(B) < \infty$ . As  $\tilde{\mu}$  is  $\mathcal{K}_\delta$ -regular, we can find a  $\mathcal{K}_\delta$ -set  $\bar{K}$  such that  $\bar{K} \subset B$  and  $a < \tilde{\mu}(\bar{K}) < \infty$ . Then  $\tilde{\mu}(\bar{K}) = \tilde{\mu}_1(\bar{K})$  which proves (3.4).

For any  $A_0 \in \mathcal{A}$  with  $\mu(A_0) < \infty$ , we have  $\infty > \mu(A_0) = \tilde{\mu}(A_0) = \tilde{\mu}_1(A_0)$  by (3.3). This implies, for any  $\mathcal{A}$ -set  $A$ ,  $\tilde{\mu}_1(A) \leq \tilde{\mu}(A) \leq \mu(A) = \sup\{\mu(A_0) : A_0 \in \mathcal{A}, A_0 \subset A, \mu(A_0) < \infty\} = \sup\{\tilde{\mu}_1(A_0) : A_0 \in \mathcal{A}, A_0 \subset A, \mu(A_0) < \infty\} \leq \tilde{\mu}_1(A)$ . Thus  $\tilde{\mu}_1$  is a semifinite,  $\mathcal{K}_\delta$ -regular measure extension of  $\mu$ . ■

**3.5. PROPOSITION.** Assume that  $\mathcal{K} \subset \mathcal{L}$  where  $\mathcal{L}$  is sequentially dominated by  $\mathcal{A}$ .

(a) If  $\mu$  is a (not necessarily semifinite) measure on  $\mathcal{A}$  such that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ , then  $\mu$  can be extended to a measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$ .

(b) If  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  for every finite measure  $\mu$  on  $\mathcal{A}$ , then every measure on  $\mathcal{A}$  can be extended to a measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$ .

*Proof.* Let  $\mu$  be an arbitrary measure on  $\mathcal{A}$ . For any  $A \in \mathcal{A}$ , put  $\nu(A) := \sup\{\mu(A_0) : A_0 \in \mathcal{A}, A_0 \subset A \text{ and } \mu(A_0) < \infty\}$  and  $\tau(A) := \sup\{\mu(B) - \nu(B) : B \in \mathcal{A}, B \subset A \text{ and } \nu(B) < \infty\}$ . It is easy to verify that

$$\mu = \nu + \tau, \quad \text{where } \nu [\tau] \text{ is a semifinite } [\{0, \infty\}\text{-valued}] \text{ measure on } \mathcal{A}. \quad (3.5)$$

In particular,  $\tau$  can be extended to a measure  $\tau_1$  on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$  (see [3, 1.3]).

Ad (a). Let  $\mu$  be a measure on  $\mathcal{A}$  such that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ . Decompose  $\mu$  according to (3.5). Then it is easy to see that  $\mathcal{K}$   $\nu$ -approximates  $\mathcal{A}$ . Thus we infer from 3.4(b) that  $\nu$  can be extended to a measure  $\nu_1$  on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$ . Hence  $\nu_1 + \tau_1$  is the desired measure extension of  $\mu$ .

Ad (b). Assume that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  for every finite measure  $\mu$  on  $\mathcal{A}$ . This implies that  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$  for every semifinite measure  $\mu$  on  $\mathcal{A}$ . Using the decomposition (3.5), our claim now follows from 3.4(b). ■

Under the additional assumption  $\mathcal{K} \subset \mathcal{A}$ , Theorem 3.4 can be strengthened in the following way.

3.6. THEOREM. *Assume that  $\mathcal{K} \subset \mathcal{A} \cap \mathcal{L}$ . Then we have:*

(a) *Every semifinite,  $\mathcal{K}$ -regular content  $\mu$  on  $\mathcal{A}$  can be extended to a semifinite,  $\mathcal{L}$ -regular content on  $\alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{K}) \cup \mathcal{F}(\mathcal{L}))$ . The extension is unique on every algebra  $\mathcal{A}'$  with  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{K}) \cup \mathcal{F}(\mathcal{L}))$  if  $\mu$  is finite and  $\alpha(\mathcal{A} \cup \mathcal{F}(\mathcal{K}))$  separates  $\mathcal{L}$ .*

(b) *If, in addition,  $\mathcal{L}$  is sequentially dominated by  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$ , then every semifinite,  $\mathcal{K}$ -regular measure  $\mu$  on  $\mathcal{A}$  can be extended to a semifinite,  $\mathcal{L}_\delta$ -regular measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$ . The extension is unique on every  $\sigma$ -algebra  $\mathcal{A}'$  with  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$  if  $\mu$  is  $\sigma$ -finite and  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$  separates  $\mathcal{L}_\delta$ .*

*Proof.* We only prove (b), since the (simpler) proof of (a) can be performed in the same way. Define  $\mathcal{C} := \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$  and  $\mathcal{D} := \sigma(\mathcal{C} \cup \mathcal{F}(\mathcal{L}_\delta)) = \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$ . Let  $\mu$  be a semifinite,  $\mathcal{K}$ -regular measure on  $\mathcal{A}$ . Set  $\lambda := \mu|_{\mathcal{K}}$ . Then  $\lambda \in N(\mathcal{K})$  is tight, semifinite, and  $\sigma$ -smooth at  $\emptyset$ .

Now we let  $A \in \mathcal{A}$  be given. For any  $K \in \mathcal{K}$ , we have  $\lambda(K) = \mu(K) = \mu(K \cap A) + \mu(K - A) = \lambda_*(K \cap A) + \lambda_*(K - A)$  which implies  $A \in \mathcal{M}(\lambda_*)$  by [2, 1.1d]. Thus we have  $\mathcal{A} \subset \mathcal{M}(\lambda_*)$ , and from 1.3(a)–(c) we obtain  $\mathcal{C} \subset \mathcal{M}(\lambda^s)$ , where  $\nu := \lambda^s|_{\mathcal{C}}$  is a semifinite,  $\mathcal{K}_\delta$ -regular measure which extends  $\mu = \lambda_*|_{\mathcal{A}}$ . Since  $\mathcal{L}$  is sequentially dominated by  $\mathcal{C}$ , so is  $\mathcal{L}_\delta$ . In addition, we have  $\mathcal{K}_\delta \subset \mathcal{L}_\delta$ . Thus we infer from 3.4(b) that  $\nu$  can be extended to a semifinite,  $\mathcal{L}_\delta$ -regular measure on  $\mathcal{D}$ .

In order to prove the uniqueness statement, let  $\mathcal{A}'$  be any  $\sigma$ -algebra satisfying  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \mathcal{D}$ . Assume that  $\mathcal{C}$  separates  $\mathcal{L}_\delta$  and that  $\mu$  is finite. (The  $\sigma$ -finite case can be reduced to the finite case as in the proof of 3.1(b).) Let  $\mu_1, \mu_2$  be  $\mathcal{L}_\delta$ -regular measures on  $\mathcal{A}'$  that extend  $\mu$ . Put  $\rho := \mu_1|_{\mathcal{L}_\delta}$  and  $\tau := \mu_2|_{\mathcal{L}_\delta}$ .  $\rho$  and  $\tau$  are finite, tight set functions on  $\mathcal{L}_\delta$  being  $\sigma$ -smooth at  $\emptyset$ .

For any  $Q \subset X$ , we have  $\lambda^s(Q) = \sup\{\lambda^*(\bar{K}): \bar{K} \in \mathcal{K}_\delta, \bar{K} \subset Q\} \leq \sup\{\mu_1(\bar{K}): \bar{K} \in \mathcal{K}_\delta, \bar{K} \subset Q\} \leq \sup\{\mu_1(\bar{L}): \bar{L} \in \mathcal{L}_\delta, \bar{L} \subset Q\} = \rho_*(Q)$ , and  $\lambda^s(X) = \mu(X) = \mu_1(X) = \rho_*(X)$ . Thus we have

$$\lambda^s \leq \rho_* \quad \text{and} \quad \lambda^s(X) = \rho_*(X). \quad (3.6)$$

Furthermore, we have

$$\mathcal{F}(\mathcal{K}_\delta) \subset \mathcal{M}(\rho_*). \quad (3.7)$$

If  $F \in \mathcal{F}(\mathcal{K}_\delta)$ , then  $F \in \mathcal{M}(\lambda^s)$  by 1.3(a), hence  $\rho_*(F) + \rho_*(X - F) \leq \rho_*(X) = \lambda^s(X) = \lambda^s(F) + \lambda^s(X - F) \leq \rho_*(F) + \rho_*(X - F)$  by (3.6). Therefore we have  $\rho_*(F) + \rho_*(X - F) = \rho_*(X)$  and so  $F \in \mathcal{M}(\rho_*)$  by 1.2. This proves (3.7).

Moreover, we have  $\mathcal{F}(\mathcal{L}_\delta) \subset \mathcal{M}(\rho_*)$  and  $\mathcal{A} \subset \mathcal{M}(\rho_*)$  by 1.1 and 1.2, respectively. Since  $\mathcal{M}(\rho_*)$  is a  $\sigma$ -algebra (see 1.3(a) and (d)), we obtain  $\mathcal{D} \subset \mathcal{M}(\rho_*)$ . As  $\lambda^s$  and  $\rho_*$  are finite measures on  $\mathcal{C}$ , (3.6) implies  $\lambda^s|_{\mathcal{C}} = \rho_*|_{\mathcal{C}}$ . In the same way, we can deduce  $\lambda^s|_{\mathcal{C}} = \tau_*|_{\mathcal{C}}$  and  $\mathcal{D} \subset \mathcal{M}(\tau_*)$ . Thus  $\rho_*|_{\mathcal{D}}$  and  $\tau_*|_{\mathcal{D}}$  are finite,  $\mathcal{L}_\delta$ -regular measures (see 1.3) which coincide on  $\mathcal{C}$ . Hence 3.1(b) implies  $\rho_*|_{\mathcal{D}} = \tau_*|_{\mathcal{D}}$ , in particular,  $\mu_1 = \mu_2$ . ■

**3.7. COROLLARY.** Assume that  $\mathcal{K} \subset \mathcal{L}$ , where  $\mathcal{L}$  is a semicompact lattice. Furthermore, let  $\mu$  be a semifinite content on  $\mathcal{A}$  such that  $\mathcal{K} \mu$ -approximates  $\mathcal{A}$ .

(a) Then  $\mu$  can be extended to a semifinite,  $\mathcal{L}_\delta$ -regular measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$ . The extension is unique on every  $\sigma$ -algebra  $\mathcal{A}'$  with  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{L}_\delta))$  if  $\mu$  is  $\sigma$ -finite and  $\mathcal{A}$  separates  $\mathcal{L}_\delta$ .

(b) If, in addition,  $\mathcal{K} \subset \mathcal{A}$  (i.e.,  $\mu$  is  $\mathcal{K}$ -regular), then  $\mu$  can be extended to a semifinite,  $\mathcal{L}_\delta$ -regular measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$ . The extension is unique on every  $\sigma$ -algebra  $\mathcal{A}'$  with  $\mathcal{A} \cup \mathcal{L} \subset \mathcal{A}' \subset \sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$  if  $\mu$  is  $\sigma$ -finite and  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$  separates  $\mathcal{L}_\delta$ .

*Proof.* Since  $\mathcal{L}$  is semicompact, so is  $\mathcal{K}$ . Thus  $\mu$  is  $\sigma$ -additive by 3.3(c). On the other hand, the semicompactness of  $\mathcal{L}$  implies that  $\mathcal{L}$  is sequentially dominated by  $\mathcal{A}$ . Now the assertion follows from 3.4(b) and 3.6(b), respectively. ■

Theorem 3.6 is a common generalization of [3, 3.2 and 3.3(a); 5, 2.1; and 18, 2.2]. Since any  $\mathcal{K}$ -regular content [measure] on a ring [ $\sigma$ -ring]  $\mathcal{R}$  can be uniquely extended to a  $\mathcal{K}$ -regular content [measure] on  $\alpha(\mathcal{R})$  [ $\sigma(\mathcal{R})$ ], we also obtain from 3.6(a) [3.7(b)] Satz 3.1 [Satz 4.5] of [11]. The following result is an immediate consequence of 3.7(b).

**3.8. COROLLARY.** Assume that  $\mathcal{K} \subset \mathcal{L} \subset \mathcal{F}(\mathcal{K}_\delta)$  where  $\mathcal{K}$  is a semicom-

compact lattice. If  $\mathcal{A} \subset \sigma(\mathcal{L})$ , then every semifinite,  $\mathcal{K} \cap \mathcal{A}$ -regular content on  $\mathcal{A}$  can be extended to a semifinite,  $\mathcal{K}_\delta$ -regular measure on  $\sigma(\mathcal{L})$ .

Corollary 3.8 is a generalization of Henry's extension theorem (see [17, Theorem 16, p. 51]) which one obtains from 3.8 if  $\mathcal{K}$ ,  $\mathcal{L}$  are the lattices of compact, respectively closed, subsets of a Hausdorff space  $X$ .

The following proposition can be derived from 3.6(b) in the same way as 3.5 has been deduced from 3.4(b).

**3.9. PROPOSITION.** Assume that  $\mathcal{K} \subset \mathcal{A} \cap \mathcal{L}$ , where  $\mathcal{L}$  is sequentially dominated by  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta))$ .

(a) If  $\mu$  is a (not necessarily semifinite)  $\mathcal{K}$ -regular measure on  $\mathcal{A}$ , then  $\mu$  can be extended to a measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$ .

(b) If every finite measure on  $\mathcal{A}$  is  $\mathcal{K}$ -regular, then every measure on  $\mathcal{A}$  can be extended to a measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}_\delta) \cup \mathcal{F}(\mathcal{L}_\delta))$ .

Further applications of 3.6 to extension problems in abstract and topological measure theory can be found in [3, Sect. 3]. The following two simple examples show that the assumption  $\mathcal{K} \subset \mathcal{A}$  is essential for the validity of 3.6.

**3.10. EXAMPLES.** (a) Let  $X$  be a set with cardinality  $\aleph_1$  and put  $\mathcal{A} := \{A \subset X : A \text{ or } X - A \text{ is countable}\}$ . For any  $\mathcal{A}$ -set  $A$ , define  $\mu(A) = 0$  or 1 according as  $A$  or  $X - A$  is countable. Furthermore, let  $\mathcal{K} = \mathcal{L} = \mathcal{P}(X)$ . Then  $(X, \mathcal{A}, \mu)$  is a probability space and  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ . In view of Ulam's theorem (see [13, Theorem 5.6]),  $\mu$  cannot be extended to a measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K})) = \mathcal{P}(X)$ .

(b) Let  $X$  be a set consisting of the three elements  $x, y, z$ . Put  $\mathcal{K} = \mathcal{L} = \mathcal{P}(X)$ . Then  $\mathcal{K}$  is a semicompact  $\delta$ -lattice. Furthermore, let  $A := \{x, y\}$  and  $\mathcal{A} := \{\emptyset, X, A, X - A\}$ . If we set  $\mu(\emptyset) = \mu(X - A) = 0$  and  $\mu(A) = \mu(X) = 1$ , then  $(X, \mathcal{A}, \mu)$  is a probability space and  $\mathcal{K}$   $\mu$ -approximates  $\mathcal{A}$ . However, the Dirac measures pertaining to the points  $x$  and  $y$ , respectively, are different ( $\mathcal{K}$ -regular) measures on  $\mathcal{P}(X)$  which extend  $\mu$ .

We now give an example of a semicompact  $\delta$ -lattice  $\mathcal{K}$  which does not  $v|_{\mathcal{A}}$ -approximate  $\mathcal{A}$  where  $v$  is a finite,  $\mathcal{K}$ -regular measure on  $\sigma(\mathcal{A} \cup \mathcal{F}(\mathcal{K}))$ . This means that the existence statements in Theorem 3.4 cannot be reversed.

**3.11. EXAMPLE.** Let  $X$  be an infinite Hausdorff space such that every real-valued continuous function on  $X$  is constant (cf. [8]). Then the Baire  $\sigma$ -algebra of  $X$  is trivial, i.e.,  $\mathcal{B}_0(X) = \{\emptyset, X\}$ . Fix some point  $x_0 \in X$  and

denote by  $\nu$  the Dirac measure pertaining to  $x_0$ , i.e., for any  $B \subset X$ ,  $\nu(B) = 1$  or 0 according as  $x_0 \in B$  or  $x_0 \notin B$ . Obviously  $\nu$  is  $\mathcal{K}(X)$ -regular. However, if  $\mu$  denotes the restriction of  $\nu$  onto  $\mathcal{B}_0(X)$ , then  $\mathcal{K}(X)$  does not  $\mu$ -approximate  $\mathcal{B}_0(X)$ , since the condition  $\mu(X) = \sup\{\mu_*(K): K \in \mathcal{K}(X)\}$  is violated. On the other hand,  $\mu$  satisfies the following weaker condition

$$\mu(X) = \sup\{\mu^*(K): K \in \mathcal{K}(X)\}. \quad (3.8)$$

According to [19], a finite Baire measure  $\mu$  on an arbitrary topological space  $X$  which satisfies (3.8) is said to be *tight*. As a consequence of the following measure extension theorem, it will be shown in 3.16 that any tight Baire measure on a Hausdorff space  $X$  can be extended to a Radon measure on  $X$ .

**3.12. THEOREM.** *Assume that the lattices  $\mathcal{K}, \mathcal{L}$  satisfy the following conditions:*

- (a)  $\mathcal{K}$  is a  $\delta$ -lattice.
- (b) For any  $K \in \mathcal{K}$ , the set  $\{L \in \mathcal{L}: K \subset L\}$  is nonempty.
- (c) If  $K \in \mathcal{K}$  and  $L \in \mathcal{L}$ , then  $K - L \in \mathcal{K}$  and  $L - K \in \mathcal{L}$ .
- (d)  $\mathcal{L}$  separates  $\mathcal{K}$ .

Furthermore, let  $\mu$  be a finite measure defined on a sub- $\sigma$ -algebra  $\mathcal{A}$  of  $\sigma(\mathcal{F}(\mathcal{K}))$  such that

$$\mu(X) = \sup\{\mu^*(K): K \in \mathcal{K}\}; \quad (3.9)$$

$$\mu^*(K) = \inf\{\mu^*(L): K \subset L \in \mathcal{L}\} \quad \text{for any } K \in \mathcal{K}; \quad (3.10)$$

$$\mu^*|_{\mathcal{K}} \text{ is } \sigma\text{-smooth at } \emptyset. \quad (3.11)$$

Then  $\mu$  can be extended to a  $\mathcal{K}$ -regular measure on  $\sigma(\mathcal{F}(\mathcal{K}))$ . The extension is unique provided that  $\mathcal{A}$  separates  $\mathcal{K}$ .

*Proof.* Choose an increasing sequence  $(K_n)$  in  $\mathcal{K}$  such that  $\mu(X) = \sup_n \mu^*(K_n)$ . As  $\nu := \mu^*|_{\mathcal{K}}$  satisfies the conditions (B1)–(B5) of [1], there exists, by [1, 3.7], a  $\mathcal{K}$ -regular measure  $\rho$  on  $\sigma(\mathcal{F}(\mathcal{K}))$  such that  $\rho|_{\mathcal{K}} \leq \nu$  and  $\rho(K_n) = \nu(K_n)$  for all  $n \in N$ . For any  $\mathcal{A}$ -set  $A$ , we thus obtain  $\rho(A) = \sup\{\rho(K): K \in \mathcal{K}, K \subset A\} \leq \sup\{\nu(K): K \in \mathcal{K}, K \subset A\} \leq \mu^*(A) = \mu(A)$ . On the other hand, we have  $\rho(X) \geq \sup_n \rho(K_n) = \sup_n \nu(K_n) = \mu(X)$ , hence  $\rho(X) = \mu(X)$  and so  $\rho|_{\mathcal{A}} = \mu$ . The uniqueness statement follows from 3.1. ■

Now we shall apply 3.12 to some topological situations. Our first application is concerned with the extension of a set function to a Baire measure. Note that any finite Baire measure on a topological space  $X$  is  $\mathcal{L}(X)$ -regular (see [19, Theorem 18, Part I]).

3.13. COROLLARY. *Let  $X$  be a topological space,  $\mathcal{A}$  a sub- $\sigma$ -algebra of  $\mathcal{B}_0(X)$  and  $\mu$  a finite measure on  $\mathcal{A}$  such that  $\mu^*|_{\mathcal{L}(X)}$  is  $\sigma$ -smooth from above (i.e.,  $\mu^*(\bigcap_n Z_n) = \inf_n \mu^*(Z_n)$  for every decreasing sequence  $(Z_n)$  in  $\mathcal{L}(X)$ ). Then  $\mu$  can be extended to a Baire measure on  $X$ . The extension is unique if  $\mathcal{A}$  separates  $\mathcal{L}(X)$ .*

*Proof.* Put  $\mathcal{K} := \mathcal{L}(X)$  and  $\mathcal{L} := \mathcal{U}(X)$ . Then the conditions (a)–(d) of 3.12 are satisfied (cf. [1, 3.15]). In addition, we have  $\mathcal{F}(\mathcal{K}) = \mathcal{K}$ , hence  $\sigma(\mathcal{F}(\mathcal{K})) = \mathcal{B}_0(X)$ . Since  $\mu$  trivially satisfies the conditions (3.9) and (3.11), it remains to verify (3.10). Let  $K \in \mathcal{L}(X)$  be given. Then there exist two decreasing sequences  $(K_n) \subset \mathcal{L}(X)$  and  $(L_n) \subset \mathcal{U}(X)$  such that  $K = \bigcap_n L_n = \bigcap_n K_n$  and  $L_n \subset K_n$  for all  $n \in \mathbb{N}$ . This implies  $\mu^*(K) = \inf_n \mu^*(K_n) \geq \inf_n \mu^*(L_n) \geq \inf\{\mu^*(L) : K \subset L \in \mathcal{U}(X)\} \geq \mu^*(K)$ . ■

Our next applications of 3.12 are concerned with Borel extensions.

3.14. COROLLARY. *Let  $X$  be a normal topological space,  $\mathcal{A}$  a sub- $\sigma$ -algebra of  $\mathcal{B}(X)$  and  $\mu$  a finite measure on  $\mathcal{A}$  such that*

- (i)  $\mu^*(F) = \inf\{\mu^*(G) : F \subset G \in \mathcal{G}(X)\}$  for all  $F \in \mathcal{F}(X)$ ;
- (ii)  $\mu^*|_{\mathcal{F}(X)}$  is  $\sigma$ -smooth at  $\emptyset$ .

*Then  $\mu$  can be extended to an  $\mathcal{F}(X)$ -regular Borel measure on  $X$ . The extension is unique if  $\mathcal{A}$  separates  $\mathcal{F}(X)$ .*

*Proof.* Apply 3.12 to  $\mathcal{K} := \mathcal{F}(X)$  and  $\mathcal{L} := \mathcal{G}(X)$ . ■

The existence part of the following corollary was proved by different methods in [7, Proposition 4.7(c)].

3.15. COROLLARY. *Let  $X$  be a Hausdorff space and let  $\mu$  be a finite measure defined on a sub- $\sigma$ -algebra  $\mathcal{A}$  of  $\mathcal{B}(X)$ . If  $\mu$  satisfies the following two conditions*

- (i)  $\mu(X) = \sup\{\mu^*(K) : K \in \mathcal{K}(X)\}$ ,
- (ii)  $\mu^*(K) = \inf\{\mu^*(G) : K \subset G \in \mathcal{G}(X)\}$  for all  $K \in \mathcal{K}(X)$ ,

*then  $\mu$  can be extended to a Radon measure on  $X$ . The extension is unique if  $\mathcal{A}$  separates  $\mathcal{K}(X)$ .*

*Proof.* Our claim follows from 3.12 with  $\mathcal{K} := \mathcal{K}(X)$  and  $\mathcal{L} := \mathcal{G}(X)$ . Note that in this case condition (3.11) is automatically satisfied. ■

In the special case  $\mathcal{A} = \mathcal{B}_0(X)$  we obtain from 3.15

3.16. COROLLARY. *Every finite tight Baire measure on a Hausdorff space  $X$  can be extended to a Radon measure on  $X$ . The extension is unique if  $X$  is*

completely Hausdorff (i.e., the continuous real-valued functions separate the points of  $X$ ).

*Proof.* Condition (ii) of the preceding corollary is satisfied since we have  $\mu(B) = \inf\{\mu(U) : B \subset U \in \mathcal{U}(X)\}$ ,  $B \in \mathcal{B}_0(X)$ , for every finite Baire measure  $\mu$  on  $X$ . ■

#### ACKNOWLEDGMENTS

The author is very much indebted to Jürgen Kindler for several helpful remarks which, in particular, led to a simplified proof of 2.2.

#### REFERENCES

1. W. ADAMSKI, Capacitylike set functions and upper envelopes of measures, *Math. Ann.* **229** (1977), 237–244.
2. W. ADAMSKI, Tight set functions and essential measure, *Lecture Notes in Math.* Vol. 945, pp. 1–14, Springer-Verlag, Berlin/Heidelberg/New York, 1982.
3. W. ADAMSKI, Extensions of tight set functions with applications in topological measure theory, *Trans. Amer. Math. Soc.* **283** (1984), 353–368.
4. B. ANGER AND J. LEMBCKE, Hahn–Banach type theorems for hypolinear functionals, *Math. Ann.* **209** (1974), 127–151.
5. G. BACHMAN AND A. SULTAN, On regular extensions of measures, *Pacific J. Math.* **86** (1980), 389–395.
6. F. DELBAEN, Convex games and extreme points, *J. Math. Anal. Appl.* **45** (1974), 210–233.
7. S. GRAF, Measurable weak selections, *Lecture Notes in Math.* Vol. 794, pp. 117–140, Springer-Verlag, Berlin/Heidelberg/New York, 1980.
8. E. HEWITT, On two problems of Urysohn, *Ann. of Math.* (2) **47** (1946), 503–509.
9. Y. KANNAI, Countably additive measures in cores of games, *J. Math. Anal. Appl.* **27** (1969), 227–240.
10. J. L. KELLEY, M. K. NAYAK, AND T. P. SRINIVASAN, Pre-measures on lattices of sets, II, in “Vector and Operator Valued Measures and Applications” (D. H. Tucker and H. B. Maynard, Eds.), pp. 155–164, Academic Press, New York, 1973.
11. J. LEMBCKE, Konservative Abbildungen und Fortsetzung regulärer Maße, *Z. Wahrsch. Verw. Gebiete* **15** (1970), 57–96.
12. E. MARCZEWSKI, On compact measures, *Fund. Math.* **40** (1953), 113–124.
13. J. C. OXTOPY, “Measure and Category,” 2nd ed., Springer-Verlag, New York/Heidelberg/Berlin, 1980.
14. J. K. PACHL, Disintegration and compact measures, *Math. Scand.* **43** (1978), 157–168.
15. J. PFANZAGL AND W. PIERLO, “Compact Systems of Sets,” *Lecture Notes in Math.* Vol. 16, Springer-Verlag, Berlin/Heidelberg/New York, 1966.
16. D. SCHMEIDLER, Cores of exact games I, *J. Math. Anal. Appl.* **40** (1972), 214–225.
17. L. SCHWARTZ, “Radon Measures on Arbitrary Topological Spaces and Cylindrical Measures,” Oxford Univ. Press, Bombay, 1973.
18. M. SZETO, On maximal measures with respect to a lattice, in “Measure Theory and its Applications” (G. A. Goldin and R. F. Wheeler, Eds.), pp. 277–282, Northern Illinois Univ., De Kalb, IL, 1981.
19. V. S. VARADARAJAN, Measures on topological spaces, *Amer. Math. Soc. Transl.* **48** (1965), 161–228.